

[54] **METHOD FOR PRODUCING EGGS
CONTAINING A HIGH AMOUNT OF
IODINATED AMINO ACIDS**

[75] Inventors: **Tadashi Ishikawa**, Sagamihara;
Hiroshi Kamimae, Yokohama, both
of Japan

[73] Assignee: **Nihon Nousan Kougyou Co., Ltd.**,
Yokohama, Japan

[21] Appl. No.: **922,581**

[22] Filed: **Jul. 6, 1978**

[51] Int. Cl.² **A61K 33/18**

[52] U.S. Cl. **424/150; 424/319**

[58] Field of Search **424/150, 319**

[56] **References Cited**

U.S. PATENT DOCUMENTS

3,526,508 9/1970 Andelfinger et al. 424/150

OTHER PUBLICATIONS

Chemical Abstracts, vol. 82, (1975), p. 123663u.

Chemical Abstracts, vol. 70, (1969), p. 9632k.

Primary Examiner—V. D. Turner

Attorney, Agent, or Firm—Haseltine, Lake & Waters

[57]

ABSTRACT

A method for producing eggs that contain a high amount (i.e., 7 ppm and over) of iodinated amino acids. Iodinated amino acids are transferred to eggs from live birds such as chickens, ducks and other poultry that are regularly fed with a specially prepared feed containing more than 230 mg. of calcium iodate per kg. of feed. Said eggs are effective in preventing and curing hypercholesteremia.

1 Claim, No Drawings